

AMPROL® 25%

...offers both preventative and treatment programs for coccidiosis.

AMPROL 25% offers the flexibility of treatment and prevention programs for a wide variety of poultry species in a medicated feed. With a number of different medication levels from which to choose, a solution to your coccidiosis problem can be obtained.

DIRECTIONS FOR USE:

Broilers:

As an aid in prevention, use a medicated feed containing 0.0125% amprolium on a continuous basis. Where severe coccidiosis conditions exist use a feed containing amprolium at the rate of 0.025%. In situations where *E. tenella* is the major problem use an amprolium medicated feed at a rate of 0.008%-0.0125%.

Replacements:

For use as an aid in the development of immunity or the prevention of coccidiosis. Where the development of immunity is not desired, use a medicated feed containing a level of 0.0125%-0.025% amprolium on a continuous basis from day one to the onset of production. In situations where immunity is desired use a medicated feed containing a level of 0.004%-0.0125% fed continuously until the onset of production. Selection of the correct level to be used should be based upon the comparative hazard of infection with cecal and intestinal species of *coccidia*. Higher levels will interfere with the development of immunity to *E. tenella*. The following are suggested feeding levels of amprolium for various conditions of coccidial exposure in replacements.

<u>Weeks of Age</u>	<u>Severe Exposure</u>	<u>Moderate Exposure</u>	<u>Slight Exposure</u>
0-5 weeks	0.0125%	0.008%-0.0125%	0.004%-0.0125%
5-8 weeks	0.008%-0.0125%	0.006%-0.0125%	0.004%-0.0125%
over 8 weeks	0.004%-0.0125%	0.004%-0.0125%	0.004%-0.0125%

* AMPROL, The Enteric shield, Building Blocks to Performance, and Performance Builder Logo are all trademarks of Merial, ©2004 Merial Select, Gainesville, GA. All rights reserved.



Laying Hens:

For treatment, use a medicated feed containing 0.025% amprolium for two weeks in severe outbreaks or 0.0125% for two weeks in moderate outbreaks of coccidiosis. In situations where prevention is the objective, use a 0.0125% medicated feed continuously.

Turkeys:

For prevention, use a medicated feed containing 0.0125% amprolium continuously.

Pheasants:

For the prevention of coccidiosis in growing pheasants caused by *E. colchici*, *E. duodenalis*, and *E. phasiani*, a medicated feed containing 0.0175% amprolium should be used continuously.

CAUTIONS:

Use as the sole source of amprolium. Do not change the litter while using this feed unless absolutely necessary. If losses exceed 0.5% in a 2-day period, obtain an accurate diagnosis, and follow the instructions of your veterinarian or poultry pathologist. Losses may result from intercurrent disease or other conditions affecting drug intake, which can contribute to the virulence of coccidiosis under field conditions. In replacement flocks the grower must expect that excessive exposure to one or more species of *coccidia* may overwhelm the drug in some flocks and prompt treatment will be required. Fertility, hatchability, and other reproductive data are not available for amprolium use in breeding pheasants. Do not use in feeds containing bentonite.

MIXING DIRECTIONS:

AMPROL 25% should be thoroughly and evenly mixed in accordance with current good manufacturing practice for medicated feed. The table below shows the amount of AMPROL 25% to be used in each ton (2000 lb.) of feed to obtain the desired level of amprolium.

<u>Feeding Level of Amprolium</u>	<u>Use this amount of AMPROL® 25% Per Ton</u>
0.004%	5 oz.
0.006%	8 oz.
0.008%	10 oz.
0.0125%	1 lb.
0.0175%	1 lb. 6 1/2 oz.
0.025%	2 lb.

*For questions or concerns about the AMPROL product line please call 1-877-4AMPROL (1-877-426-7765)



MERIAL SELECT
P.O. Drawer 2497
Gainesville, GA 30503
770-536-8787
770-534-0998 fax