

# ANIT-COX



*Specification Sheet*

TYPE "B" MEDICATED ARTICLE

www.asp-inc.com

## INDICATIONS

A type "B" medicated article specifically formulated for uniform dispersion in cattle and goat feeds.

## ADVANTAGES

- For the prevention of coccidiosis in ruminating and non-ruminating calves and cattle caused by Eimeria Bovis and E. Zurni.
- Type "B" medicated article does not require 1900 registration,
- Convenient, free flowing formulation.

## INGREDIENTS

### Guaranteed Analysis:

Decoquinatate.....2.27 gm/lb

Rice hulls, Calcium carbonate, Soybean oil, Corn meal, Lecithin.

## CAUTIONS

Do not feed to cows producing milk for food.  
Do not mix in feeds containing bentonite.  
For use is animal feeds only.  
Store in a cool, dry place.  
Keep out of reach of children.

## PACKAGING

25 lb. bag

Specially designed multi-wall bag contains 2 exterior walls of paper for protection, an inner-poly liner for moisture protection followed by an interior paper liner.

## DIRECTIONS FOR USE

### Mixing Directions:

**Cattle:** Feed at the rate of 22.7 mg per 100 pounds of bodyweight per day for 28 days.

**Goats:** For young goats, feed at the rate of 22.7 mg per 100 pounds of bodyweight per day for 28 days.

Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be hazard. Coccidiostats are not indicated for use in adult animals due to continuous exposure.

To feed this level: Premix 10 pounds of ANTI-COX with 90 pounds of grain. The resulting premix should be topdressed or mixed into the daily ration at 1/10 lb. per 100 lbs. bodyweight.