

Indications

Chickens: As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. Acervulina*, *E. brunetti*, *E. Mivati* and *E. maxima* in broiler chickens and replacement chickens intended for use as caged layers.

Turkeys: For the prevention of coccidiosis in growing turkeys caused by *Eimeria adenoeides*, *E. Meleagrimitis* and *E. Gallopavonis*.

Quail: For the prevention of coccidiosis in Bobwhite quail caused by *Eimeria dispersa* and *E. lettyae*.

Use Programs

Important: Do not feed undiluted. Must be thoroughly mixed in feeds before use.

Chickens: Feed continuously as the sole ration a complete feed containing 90 to 110 grams of monensin per ton. Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.

Turkeys: Feed continuously to growing turkeys as the sole ration from one day of age a complete feed containing 54 to 90 grams of monensin per ton depending upon the severity of coccidiosis exposure. Bobwhite Quail: Feed continuously as the sole ration a complete feed containing 73 grams of monensin per ton.

Key Points

1. Effective against the six pathogenic species of chicken coccidia.
2. Effective against three pathogenic species of turkey coccidia.
3. Effective against two pathogenic species of quail coccidia.
4. Mode of action is primarily coccidiocidal.
5. Cleared for use in both broiler and layer replacement chicken feeds.
6. Cleared for use in chickens in combination with 3-Nitro (Roxarsone) and all major antibiotics.
7. Effective early in coccidial life cycle before intestinal tissue damage occurs.
8. Control of coccidiosis improves weight gains and feed conversions.

Important: For complete use directions and additional information on this product, see the reverse side of this sheet.

Coban® 60

monensin sodium
Product No. AF 1309

Type A Medicated Article

Category I Feed-Grade Anticoccidial Premix Containing 60 Grams of Monensin Per Pound (as monensin sodium)

For Use in Broiler and Replacement Chicken, Growing Turkey, and Growing Bobwhite Quail Feeds Only.

FDA STATUS: No FDA-1900 Required (Regulation §558.355). For FDA regulations on the use of Coban 60 in combination with other Type A Articles allowed in broiler chicken feed, see the current FEED ADDITIVE COMPENDIUM or consult your Elanco representative.

PACKAGE: 50-pound multiwall, flat-bottom bag

STORAGE: Store in a cool, dry place.
DATING PERIOD: 24 months



ELANCO
ANIMAL HEALTH

7M

Coban® 60

monensin sodium

Type A Medicated Article

Do Not Feed Undiluted.

Active Drug Ingredient: Monensin (as monensin sodium) 60 grams per pound.

Chickens: As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

Turkeys: For the prevention of coccidiosis in turkeys caused by *Eimeria adenoeides*, *E. meleagrimitis* and *E. gallopavonis*.

Quail: For the prevention of coccidiosis in Bobwhite quail caused by *Eimeria dispersa* and *E. lettyae*.

Important: Must be Thoroughly Mixed in Feeds Before Use.

Mixing Directions - Broiler and Replacement

Chickens: Thoroughly mix the following amounts of Coban 60 with a sufficient quantity of unmedicated feed to make one ton of complete feed to provide 90 to 110 grams of monensin per ton.

Monensin (Grams per ton)	Coban 60 (Pounds per ton of Type C Feed)
90	1.50
95	1.58
100	1.67
105	1.75
110	1.83

Feeding Directions: Feed continuously as the sole ration.



Mixing Directions -Turkeys:

Thoroughly mix the following amounts of Coban 60 with a sufficient quantity of unmedicated feed to make one ton of complete turkey feed to provide 54 through 90 grams of monensin per ton.

Monensin (Grams per ton)	Coban 60 (Pounds per ton of Type C Feed)
54	0.90
66	1.10
78	1.30
90	1.50

Feeding Directions: For turkeys, feed continuously as the sole ration. The optimum level depends upon the severity of coccidiosis exposure.

Mixing Directions - Bobwhite Quail:

Thoroughly mix the following amount of Coban 60 with a sufficient quantity of unmedicated feed to make one ton of complete quail feed to provide 73 grams of monensin per ton.

Monensin (Grams per ton)	Coban 60 (Pounds per ton of Type C Feed)
73	1.22

Feeding Directions: For Bobwhite quail, feed continuously as the sole ration.

Warning: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age.

Warning: When mixing and handling Coban 60 use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Caution: For replacement chickens intended for use as caged layers only. Do not allow horses, other equines, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses, mature turkeys and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis. In the absence of coccidiosis in broiler chickens, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain.

Store in a cool, dry place.