

Indications

Swine:

For increased rate of weight gain and improved feed efficiency.

For prevention and/or control of porcine proliferative enteropathies (ileitis).

For prevention of swine dysentery (vibriotic).

For maintaining weight gains and feed efficiency in the presence of atrophic rhinitis.

For treatment and control of swine dysentery (vibriotic) following initial medication of Tylan® Soluble in drinking water.

Beef Cattle:

For reduction of the incidence of liver abscesses caused by *Fusobacterium necrophorum* and *Actinomyces pyogenes*.

Chickens:

For increased rate of weight gain and improved feed efficiency.

Broiler and Replacement Chickens:

To aid in the control of chronic respiratory disease caused by *Mycoplasma gallisepticum*.

Laying Chickens:

For improving feed efficiency.

Important: Do not feed undiluted. Must be thoroughly mixed in feeds before use.

Key Points:

1. Tylan® increases average daily gain and improves feed efficiency as well as, or better than, competitive antibiotics used in on-farm demonstrations in swine in the 1990s — just as it did in the 70s and 80s.*
2. Tylan® prevents and/or controls ileitis.
3. Reduces the incidence of liver abscesses in beef cattle.
4. The only antibiotic cleared for combined use with Rumensin® in feedlot cattle.
5. Primarily active against Gram-positive bacteria and has significant activity against mycoplasma.
6. No withdrawal period required in swine or cattle. In chickens, a 5-day withdrawal is required only when used at 800 to 1000 grams per ton.
7. No Form FDA-1900 requirement permits easy alteration of use level, particularly in swine.

*Data available upon request from Elanco Animal Health.

Important: For complete use directions and additional information on this product, see the reverse side of this sheet.

Tylan® 10

tylosin phosphate

Type B Medicated Feed

Category 1 Feed Additive Premix
Containing 10 Grams of Tylosin Per
Pound (as tylosin phosphate)

For Use in Swine, Beef Cattle and
Chicken Feeds Only

FDA STATUS: No Feed Mill License
Required.

PACKAGE: 50-pound, multiwall,
flat-bottom bag

STORAGE: Store in a cool, dry area.

DATING PERIOD: 24 months



ELANCO
ANIMAL HEALTH

TM

* Tylan® is a trademark for Elanco's brand of tylosin tartrate.

Tylan[®] 10

tylosin phosphate

Type B Medicated Feed

Do Not Feed Undiluted

Active Drug Ingredient: Tylosin (as tylosin phosphate)
10 grams per pound

Ingredients: Rice Hulls, Calcium Carbonate and Mineral Oil.

Guaranteed Analysis:

Crude Protein, Minimum.....1.8%

Crude Fat, Minimum1.9%

Crude Fiber, Maximum35.5%

Calcium (Ca), Minimum9.5%

Calcium (Ca), Maximum11.4%

Phosphorous (P), Minimum 0.1%

This product has been formulated to contain at least 25 percent nutrient content.

For use in Swine, Beef Cattle and Chicken Feeds Only.

Important: Must be thoroughly mixed in feeds before use. Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2 percent bentonite.

Swine

**Feed Continuously As The Only Ration
(No Withdrawal Required)**

Usage	Tylan 10 per Ton of Complete Feed	Tylosin per Ton of Complete Feed
Increased rate of gain and improved feed efficiency	Starter - 2 to 10 lbs Grower - 2 to 4 lbs Finisher - 1 to 2 lbs	20 to 100 g 20 to 40 g 10 to 20 g
Prevention and/or control of ileitis	10 lbs	100 g for 21 days
Prevention of swine dysentery (vibriotic)	10 lbs ...Followed by... 4 lbs	100 g for at least 3 weeks ...Followed by... 40 g until market weight
Maintaining weight gains and feed efficiency in the presence of atrophic rhinitis	10 lbs	100 g

Beef Cattle

**Feed Continuously As The Only Ration
(No Withdrawal Required)**

Usage	Tylan 10 per Ton of Complete Feed	Tylosin per Ton of Complete Feed
For reduction of incidence of liver abscesses in beef cattle caused by <i>Fusobacterium necrophorum</i> and <i>Actinomyces pyogenes</i>	0.8 to 1.0 lb	8 to 10 g To be fed so that each animal should receive not more than 90 mg per head per day and not less than 60 mg per head per day

Chickens

**Feed Continuously As The Only Ration
(No Withdrawal Required)**

Usage	Tylan 10 per Ton of Complete Feed	Tylosin per Ton of Complete Feed
Increased rate of gain and improved feed efficiency	0.8 to 5 lbs	4 to 50 g
Laying Chickens		
Improving feed efficiency	2 to 5 lbs	20 to 50 g

TAKE TIME TO



OBSERVE LABEL DIRECTIONS