



The Leader in Trace Mineral Nutrition

# Product Information


## ZINPRO 100 ZINC METHIONINE

A nutritional feed additive for livestock and poultry. When used as a commercial feed ingredient it must be declared as zinc methionine. Association of American Feed Control Officials (AAFCO) No. 57.151. International Feed Number (IFN) 6-16-293. Zinc Methionine Complex. U.S. Patent No. 4,021,569, 4,764,633 and 5,430,164. Canada Reg. No. 990191.

### TYPICAL ANALYSIS:

Zinc	10.0%
Methionine	20.0%
Protein	11.8%
Fat	0.0%
Fiber	15.0%
Ash	20.0%
Salt	1.2%

### MOLECULAR STRUCTURE:



### PHYSICAL DESCRIPTION:

A light brown granular powder with characteristic methionine-like odor. ZINPRO 100 weighs approximately 35 lb./cu.ft. and is packaged in 50 lb. multiwall bags.

### FEEDING INSTRUCTIONS:

\*SWINE: Add 0.8 lb.(360 grams) per ton of complete ration.

LAYING HENS, BROILERS AND TURKEYS: Add 0.8 lb. (360 grams) per ton of complete ration.

DAIRY CATTLE: Feed 3.6 grams per head daily, or 1 lb. per 125 head daily.

BEEF CATTLE: Feed 3.6 grams per head daily, or 1 lb. per 125 head daily.

HORSES: Feed 3.6 grams per head daily.

\*SHEEP: Feed 0.4 grams per head daily, or 1 lb. per 1,250 head daily.

\*Double recommended levels during periods of reduced feed intake.

### TOXICITY AND SAFE HANDLING:

When correctly used, there is no toxicity hazard in the use of ZINPRO 100.