



The Leader in Trace Mineral Nutrition

# Product Information


## ZINPRO 40 ZINC METHIONINE

A nutritional feed additive for livestock and poultry. When used as a commercial feed ingredient it must be declared as zinc methionine. Association of American Feed Control Officials (AAFCO) No. 57.151. International Feed Number (IFN) 6-16-293. Zinc Methionine Complex. U.S. Patent No. 4,021,569, 4,764,633 and 5,430,164. Canada Reg. No. 980417.

### TYPICAL ANALYSIS:

Zinc	4.0%
Methionine	8.0%
Protein	5.4%
Fat	0.0%
Fiber	13.9%
Ash	51.5%
Calcium	8.5%
Salt	2.8%

### MOLECULAR STRUCTURE:



### PHYSICAL DESCRIPTION:

A light brown granular powder with characteristic methionine-like odor. ZINPRO 40 weighs approximately 32 lbs./cu.ft. and is packaged in 50 lb. multiwall bags.

### FEEDING INSTRUCTIONS:

\*SWINE: Add 2 lbs. (908 grams) per ton of complete ration.

LAYING HENS, BROILERS AND TURKEYS: Add 2 lbs. (908 grams) per ton of complete ration.

DAIRY CATTLE: Feed 9 grams per head daily, or 2 lbs. per 100 head daily.

BEEF CATTLE: Feed 9 grams per head daily, or 2 lbs. per 100 head daily

HORSES: Feed 9.0 grams per head daily. .

\*SHEEP: Feed 1 gram per head daily, or 1 lb. per 454 head daily.

\*Double recommended levels during periods of reduced feed intake.

### TOXICITY AND SAFE HANDLING:

When correctly used, there is no toxicity hazard in the use of ZINPRO 40.